

Gauge field divergences in the light-front

B.M.Pimentel^a, J.H.O.Sales^a and Tobias Frederico^b

^aInstituto de Física Teórica-UNESP, 01405-900 São Paulo, Brazil.

^bInstituto Tecnológico de Aeronáutica, CTA, 12228-900
São José dos Campos, Brazil.

February 8, 2008

Abstract

The Bethe-Salpeter equation for ground state of two fermions exchanging a gauge boson presents divergences in the momentum transverse, even in the ladder approximation projected in light-front. Gauge theories with light-front gauge also present the difficulty associated to the instantaneous term of the propagator of a system composed by fermions bosons-exchange interaction. We used a prescription that allowed an appropriate description of the singularity in the propagator of the gauge boson in the light-front.

1 Light-Front Dynamics: Definition

Beginning from Dirac's idea [1] of representing the dynamics of the quantum system at light-front times $x^+ = t + z$, we derive the Green's function from the covariant propagator that evolves the system from one light-front hyper-surface to another one. The light-front Green's function is the probability amplitude for an initial state at $x^+ = 0$ to evolve to a final state in the Fock-state at some x^+ , where the evolution operator is defined by the light-front Hamiltonian [2]

2 The Scalar Field Propagator

The Feynman propagator for the scalar field is

$$S(x^\mu) = \int \frac{d^4k}{(2\pi)^4} \frac{ie^{-ik^\mu x_\mu}}{k^2 - m^2 + i\varepsilon}. \quad (1)$$

and in terms of light-front variables [5], we have

$$S(x^+) = \frac{1}{2} \int \frac{dk^- dk^+ dk^\perp}{(2\pi)} \frac{ie^{-\frac{i}{2}k^- x^+}}{k^+ \left(k^- - \frac{k_\perp^2 + m^2 - i\varepsilon}{k^+}\right)}. \quad (2)$$

The Fourier transform of the single boson state propagator to the in the light-front time is giver by:

$$\tilde{S}(k^-) = \int dk^+ dk^\perp \frac{i}{k^+ \left(k^- - \frac{k_\perp^2 + m^2 - i\varepsilon}{k^+} \right)}. \quad (3)$$

3 Fermion Field

Let S_F denote fermion field propagator in covariant theory

$$S_F(x^\mu) = \int \frac{d^4 k}{(2\pi)^4} \frac{i(\not{k}_{\text{on}} + m)}{k^2 - m^2 + i\varepsilon} e^{-ik^\mu x_\mu}, \quad (4)$$

where $\not{k}_{\text{on}} = \frac{1}{2}\gamma^+ \frac{(k^\perp)^2 + m^2}{k^+} + \frac{1}{2}\gamma^- k^+ - \gamma^\perp k^\perp$. Using light-front variables in Eq.(4), we have

$$S_F(x^+) = \frac{i}{2} \int \frac{dk^- dk^+ dk^\perp}{(2\pi)} \left[\frac{\not{k}_{\text{on}} + m}{k^+ \left(k^- - \frac{k_\perp^2 + m^2 - i\varepsilon}{k^+} \right)} + \frac{\gamma^+}{2k^+} \right] e^{\frac{-i}{2} k^- x^+}. \quad (5)$$

We note that for the fermion field, light-front propagator differs from the Feynman propagator by an instantaneous propagator.

4 Gauge Boson Propagator

Let $S^{\mu\nu}$ gauge propagator,

$$S^{\mu\nu}(x^\mu) = \int \frac{d^4 k}{(2\pi)^4} \frac{ie^{-ik^\mu x_\mu}}{k^2 + i\varepsilon} \left[\frac{-nkg^{\mu\nu} + n^\mu k^\nu + n^\nu k^\mu}{nk} \right], \quad (6)$$

where we choose the light-front gauge $A^+ = 0$, $n^\mu = (1, 0, 0, -1)$ and the metric tensor is given from [2].

The light-front components (6) can be as written $S^{+-} = S^{-+} = S^{++} = S^{+\perp} = 0$ and

$$S^{--} = 4 \frac{ik^-}{k^+(k^2 + i\varepsilon)}, \quad S^{-\perp} = S^{\perp-} = 2 \frac{ik^\perp}{k^+(k^2 + i\varepsilon)}, \quad S^{\perp\perp} = -1 \frac{i}{k^2 + i\varepsilon} \quad (7)$$

5 Interaction in First Order

We consider the fermion-antifermion system in the light-front with one-gauge boson exchange ($A^+ = 0$), for which the interaction Lagrangian density is given by

$$\mathcal{L}_I = g\bar{\Psi}_1 \gamma_\mu A^\mu \Psi_1 + g\bar{\Psi}_2 \gamma_\nu A^\nu \Psi_2. \quad (8)$$

The fermion corresponds to the field Ψ with rest masses m and the exchanged gauge boson to the field A^μ with mass $\mu = 0$. The coupling constant is g .

The perturbative correction to the two-body propagator which comes from the exchange of one intermediate virtual boson, is

$$\begin{aligned}\Delta S_{g^2}(x^+) &= (ig)^2 \int d\bar{x}_1^+ d\bar{x}_2^+ S_{k'}(x^+ - \bar{x}_1^+)(\gamma_\mu) S_k(\bar{x}_1^+) \\ &\quad S^{\mu\nu}(\bar{x}_2^+ - \bar{x}_1^+) S_p(x^+ - \bar{x}_2^+)(\gamma_\nu) S_{p'}(\bar{x}_2^+).\end{aligned}\quad (9)$$

The intermediate boson propagates between the time interval $\bar{x}_2^+ - \bar{x}_1^+$. The labels in the particle propagators k and p indicates initial and k' and p' final states.

Performing the Fourier transform from x^+ to P^- and for the total kinematical momentum P^+ , which we choose positive, and $P^\perp$. The double integration in k^- is performed analytically in Eq.(10),

$$\begin{aligned}\Delta S_{g^2}(P^-) &= \frac{-(ig)^2 i}{(4\pi)^2} \int \frac{dk^- dk'^-}{k^+ k'^+ (P^+ - k'^+)(P^+ - k^+)} \\ &\quad \left\{ \frac{\not{k}'_{on} + m}{(k'^- - k'_{on} + \frac{i\varepsilon}{k'^+})} \gamma_- \frac{\not{k}_{on} + m}{(k^- - k_{on} + \frac{i\varepsilon}{k^+})} \right. \\ &\quad \frac{4(k^- - k'^-)}{(q^+)^2 \left(k^- - k'^- - q_{on}^- + \frac{i\varepsilon}{q^+}\right)} \frac{\not{p}'_{on} + m}{\left(p'^- - p'_{on} + \frac{i\varepsilon}{p'^+}\right)} \\ &\quad \gamma_- \frac{\not{p}_{on} + m}{\left(p^- - p_{on} + \frac{i\varepsilon}{p^+}\right)} + \\ &\quad + \frac{\not{k}'_{on} + m}{(k'^- - k'_{on} + \frac{i\varepsilon}{k'^+})} \\ &\quad \gamma_- \frac{\not{k}_{on} + m}{(k^- - k_{on} + \frac{i\varepsilon}{k^+})} \frac{2(k^\perp - k'^\perp)}{(q^+)^2 \left(k^- - k'^- - q_{on}^- + \frac{i\varepsilon}{q^+}\right)} \\ &\quad \frac{\not{p}'_{on} + m}{\left(p'^- - p'_{on} + \frac{i\varepsilon}{p'^+}\right)} \gamma_\perp \frac{\not{p}_{on} + m}{\left(p^- - p_{on} + \frac{i\varepsilon}{p^+}\right)} + [\gamma_\perp \rightarrow \gamma_-] + \\ &\quad \frac{\not{k}'_{on} + m}{(k'^- - k'_{on} + \frac{i\varepsilon}{k'^+})} \gamma_\perp \frac{\not{k}_{on} + m}{(k^- - k_{on} + \frac{i\varepsilon}{k^+})} \\ &\quad \frac{(-1)}{(q^+)^2 \left(k^- - k'^- - q_{on}^- + \frac{i\varepsilon}{q^+}\right)} \\ &\quad \left. \frac{\not{p}'_{on} + m}{\left(p'^- - p'_{on} + \frac{i\varepsilon}{p'^+}\right)} \gamma_\perp \frac{\not{p}_{on} + m}{\left(p^- - p_{on} + \frac{i\varepsilon}{p^+}\right)} \right\},\end{aligned}\quad (10)$$

6 Conclusion

From equation (10), we verified the existence of singularity in the components $(--)$ and $(-\perp)$ in the coordinate $q^+ = k^+ - k'^+$. We hoped to remove those singularity using the technique of displacement (δ^+) of the pole of the phase space in q^+ [6].

Acknowledgement 1 *T.F and B.M. Pimentel thank to CNPq for partial support. J.H.O. Sales is supported by FAPESP/Brazil pos-doctoral fellowship. We acknowledge discussions with J.F.Libonati and J.Messias.*

References

- [1] P. A. M. Dirac, Rev. Mod. Phys. **21**, 392 (1949).
- [2] A.Harindranath, *Light-Front Quantization and Non-Perturbative QCD*, editors J.P.Vary and F.Wölz, International Institute of Theoretical and Applied Physics (1997).
- [3] S.D.Glasek and K.G.Wilson Phys.Rev.**D49** (1994) 4214.
- [4] J.H.O. Sales, T.Frederico, B.V. Carlson and P.U.Sauer, Phys. Rev. **C63**:064003(2001).
- [5] J.H.O. Sales, T.Frederico, B.V. Carlson and P.U.Sauer, Phys. Rev. **C61**:044003(2000).
- [6] J.P.B.C.de Melo, J.H.O.Sales, T.Frederico and P.U.Sauer, Nucl. Phys.**A631** (1998) 574c.